



# Draft Genome Sequences of Five *Enterococcus* Species Isolated from the Gut of Patients with Suspected *Clostridium difficile* Infection

Eduardo Castro-Nallar,<sup>a</sup> Sandro L. Valenzuela,<sup>a</sup> Sebastián Baquedano,<sup>b</sup> Carolina Sánchez,<sup>c</sup> Fabiola Fernández,<sup>c\*</sup> Annette N. Trombert<sup>c</sup>

Center for Bioinformatics and Integrative Biology, Facultad de Ciencias Biológicas, Universidad Andrés Bello, Santiago, Chile<sup>a</sup>; Escuela de Tecnología Médica, Facultad de Ciencias, Universidad Mayor, Huechuraba, Santiago, Chile<sup>b</sup>; Centro de Genómica y Bioinformática, Facultad de Ciencias, Universidad Mayor, Santiago, Chile<sup>c</sup>

**ABSTRACT** We present draft genome sequences of five *Enterococcus* species from patients suspected of *Clostridium difficile* infection. Genome completeness was confirmed by presence of bacterial orthologs (97%). Gene searches using Hidden-Markov models revealed that the isolates harbor between seven and 11 genes involved in antibiotic resistance to tetracyclines, beta-lactams, and vancomycin.

Numerous reports link microbial compositional changes (dysbiosis) in the human gut microbiota to diverse disease states ranging from inflammatory illness to psychiatric conditions (1–4). In particular, studies in human and animal models have shown the importance of the gut microbiota's capability of providing colonization resistance against *C. difficile* (5, 6). Consequently, this announcement is part of a larger project aimed at characterizing the microbiota of individuals infected by *C. difficile* in Chile, both in terms of individual isolates and microbiota compositions.

We collected fecal samples from patients suspected of being infected by *C. difficile* that presented aqueous diarrhea associated with antimicrobial drug intake. We plated samples on blood agar medium (Merck; anaerobic conditions) and grew colonies in Brucella broth at 37°C without agitation (7). For DNA extraction, we used the Wizard genomic DNA purification kit (Promega) following the manufacturer's instructions. DNA was quantified in a fluorimeter (Qubit; Invitrogen) and its integrity was checked by agarose gel electrophoresis. We prepared sequencing libraries as in the TruSeq nano DNA LT kit (Illumina) using an average insert size of 450 bp.

We obtained between 1.7 and 2.3 million paired-end reads that we subsequently filtered to allow no undetermined bases and an average quality score per read of > Q20. We also trimmed the 5' and 3' ends to remove bases with quality scores of < Q20. The resulting reads were *de novo* assembled using a De Bruijn graph strategy as implemented in SPAdes 3.8 (8). Genome coverage was 48 to 173× (median = 109×). We interrogated the resulting contigs for evidence of contamination using the GenomePeek web server and found no evidence of contaminating DNA (9). We annotated the assembled genome sequences using the NCBI Prokaryotic Genome Annotation Pipeline (released 2013) (10). Information about the Pipeline can be found here: [https://www.ncbi.nlm.nih.gov/genome/annotation\\_prok/](https://www.ncbi.nlm.nih.gov/genome/annotation_prok/). Additionally, genome completeness was confirmed by BUSCO analysis of prokaryotic orthologs, where we found 97% of bacteria-wide orthologs present (11). Current standards suggest that genomes with > 85% orthologs present are considered high-quality genomes (reference *E. faecium* was 93% complete; accession no. NC\_017960.1) (12).

Received 28 March 2017 Accepted 28 March 2017 Published 18 May 2017

**Citation** Castro-Nallar E, Valenzuela SL, Baquedano S, Sánchez C, Fernández F, Trombert AN. 2017. Draft genome sequences of five *Enterococcus* species isolated from the gut of patients with suspected *Clostridium difficile* infection. *Genome Announc* 5: e00379-17. <https://doi.org/10.1128/genomeA.00379-17>.

**Copyright** © 2017 Castro-Nallar et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Annette N. Trombert, [annette.trombert@umayor.cl](mailto:annette.trombert@umayor.cl).

\* Present address: Fabiola Fernández, Escuela de Tecnología Médica, Facultad de Ciencias de la Salud, Universidad San Sebastián, Santiago, Chile.

**TABLE 1** Whole-genome shotgun projects accession numbers

Species	Strain	Source	Genome size (Mb)	No. of coding sequences	Accession no.	Assembly
<i>Enterococcus faecium</i>	97-19_S17	Human feces	2.7	2,915	MRYE00000000	GCA_001990575.1
<i>Enterococcus faecium</i>	97-3_S3	Human feces	2.9	3,165	MRYF00000000	GCA_001990605.1
<i>Enterococcus faecium</i>	97-6_S5	Human feces	2.5	2,698	MRYG00000000	GCA_001990565.1
<i>Enterococcus faecium</i>	97-7_S6	Human feces	3.1	3,390	MRYH00000000	GCA_001990615.1
<i>Enterococcus mundtii</i>	CGB1038-1_S1	Human feces	3.3	3,150	MSTR00000000	GCA_001990645.1

Of the five genome sequences presented here, only two were classified as known multilocus sequence types (97-19\_S17 and 97-7\_S6 as ST262 and ST822, respectively) (13). However, all strains were found to carry antibiotic resistance genes including *vanR* and *vanS* genes (ARO:3000574; ARO:3000071), *vanX*, and *vanY* genes (ARO:3000011; ARO:3000077), class B beta-lactamase (ARO:3000004), antibiotic efflux pumps (ARO:0010001), and tetracycline resistance genes (ARO:3000186; ARO:3000194; ARO:3000190; ARO:3000192; ARO:3000239; ARO:0000002), among others (14). This report highlights the need for comprehensive open genomic reference databases of human gut members to better address scientific questions regarding epidemiology, virulence and pathogenicity, and drug resistance. All five genomic sequences are compliant with the MIGS package “cultured bacteria/archaea, human-associated; version 4.0” (15).

**Accession number(s).** The whole-genome shotgun projects have been deposited in GenBank under the accession numbers provided in Table 1. The versions described in this paper are the first versions.

#### ACKNOWLEDGMENTS

E.C.-N. thanks the high-performance computing facility of George Washington University, ColonialOne, for providing data storage, support, and computing power for genomic analyses (colonialone.gwu.edu).

This work was supported by funding from INNOVA CORFO 13IDL2-23309 (A.N.T.) and from “CONICYT PAI/Concurso nacional apoyo al retorno de investigadores(as) desde el extranjero, convocatoria 2014 Folio 82140008.” (E.C.-N.).

#### REFERENCES

- Knoll RL, Forslund K, Kultima JR, Meyer CU, Kullmer U, Sunagawa S, Bork P, Gehring S. 30 December 2016. Gut microbiota differs between children with inflammatory bowel disease and healthy siblings in taxonomic and functional composition—a metagenomic analysis. *Am J Physiol Gastrointest Liver Physiol*. <https://doi.org/10.1152/ajpgi.00293.2016>.
- Castro-Nallar E, Bendall ML, Pérez-Losada M, Sabuncyan S, Severance EG, Dickerson FB, Schroeder JR, Yolken RH, Crandall KA. 2015. Composition, taxonomy and functional diversity of the oropharynx microbiome in individuals with schizophrenia and controls. *PeerJ* 3:e1140. <https://doi.org/10.7717/peerj.1140>.
- Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, Burcelin R. 2008. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes* 57:1470–1481.
- Severance EG, Yolken RH, Eaton WW. 2014. Autoimmune diseases, gastrointestinal disorders and the microbiome in schizophrenia: more than a gut feeling. *Schizophr Res* 159:14–19. <https://doi.org/10.1016/j.schres.2014.07.053>.
- Britton RA, Young VB. 2014. Role of the intestinal microbiota in resistance to colonization by *Clostridium difficile*. *Gastroenterology* 146:1547–1553. <https://doi.org/10.1053/j.gastro.2014.01.059>.
- Buffie CG, Pamer EG. 2013. Microbiota-mediated colonization resistance against intestinal pathogens. *Nat Rev Immunol* 13:790–801. <https://doi.org/10.1038/nri3535>.
- Bhardwaj S, Dhawale KBJ, Patil M, Divase S. 2013. *Enterococcus faecium* and *Enterococcus faecalis*, the nosocomial pathogens with special reference to multi-drug resistance and phenotypic characterization. *Int J Pharm Sci Pract* 2:1–10.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- McNair K, Edwards RA. 2015. GenomePeek—an online tool for prokaryotic genome and metagenome analysis. *PeerJ* 3:e1025. <https://doi.org/10.7717/peerj.1025>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614–6624.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>.
- Bradnam KR, Fass JN, Alexandrov A, Baranay P, Bechner M, Birol I, Boisvert S, Chapman JA, Chapuis G, Chikhi R, Chitsaz H, Chou WC, Corbeil J, Del Fabbro C, Docking TR, Durbin R, Earl D, Emrich S, Fedotov P, Fonseca NA, Ganapathy G, Gibbs RA, Gnerre S, Godzaridis E, Goldstein S, Haimel M, Hall G, Haussler D, Hiatt JB, Ho IY, Howard J, Hunt M, Jackman SD, Jaffe DB, Jarvis ED, Jiang H, Kazakov S, Kersey PJ, Kitzman JO, Knight JR. 2013. Assemblathon 2: Evaluating de novo methods of genome assembly in three vertebrate species. *GigaScience* 2:10. <https://doi.org/10.1186/2047-217X-2-10>.
- Buultjens AH, Lam MM, Ballard S, Monk IR, Mahony AA, Grabsch EA, Grayson ML, Pang S, Coombs GW, Robinson JO. 2016. Evolutionary

- origins of the emergent ST796 clone of vancomycin resistant *Enterococcus faecium*. PeerJ 5:e2916. <https://doi.org/10.7717/peerj.2916>.
14. Gibson MK, Forsberg KJ, Dantas G. 2015. Improved annotation of antibiotic resistance determinants reveals microbial resistomes cluster by ecology. ISME J 9:207–216. <https://doi.org/10.1038/ismej.2014.106>.
  15. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, Ashburner M, Axelrod N, Baldauf S, Ballard S, Boore J, Cochrane G, Cole J, Dawyndt P, De Vos P, DePamphilis C, Edwards R, Faruque N, Feldman R, Gilbert J, Gilna P, Glöckner FO, Goldstein P, Guralnick R, Haft D, Hancock D, Hermjakob H, Hertz-Fowler C, Hugenholtz P, Joint I, Kagan L, Kane M, Kennedy J, Kowalchuk G, Kottmann R, Kolker E. 2008. The minimum information about a genome sequence (MIGS) specification. Nat Biotechnol 26:541–547. <https://doi.org/10.1038/nbt1360>.